

# EXTENT OF RADIAL COUNTER-CURRENT HEAT EXCHANGE IN THE FOREARM MUSCLE DURING COLD EXPOSURE

Peter Tikuisis  
Michel B. Ducharme  
Defence and Civil Institute of Environmental Medicine  
North York, Canada

## INTRODUCTION

Counter-current heat exchange in the limbs is usually considered as an exchange of heat between major counter-current vessels in the axial direction. However, attention has been recently given to counter-current heat exchange between the microvessels of the vasculature away from the major axial vessels. Weinbaum et al. (1) have demonstrated that such heat exchange is 100% complete three generations before the capillary level. Heat recovery by this mechanism is considered in the radial, as opposed to the axial, direction. In this study, the extent of net heat recovery by this mechanism is estimated for the human forearm during cold exposure.

## METHOD

**1. Experimental.** Data were obtained from steady-state temperatures measured in the forearm (2) after 3 h of immersion in water at temperatures ( $T_w$ ) of 15 (N = 6), 20 (N = 5), and 30°C (N = 5). Temperatures were measured every 0.5 cm from the longitudinal axis of the forearm to the skin approximately 9 cm distal from the elbow, and from these values a volume-weighted mean was obtained. These means ( $\pm$  SE) were 18.74 (0.52), 23.41 (0.21), and 31.78 (0.17) °C with increasing  $T_w$ . Arterial blood temperatures entering the forearm were measured once before immersion and corrected for axial counter-current heat exchange; mean ( $\pm$  SE) values at steady-state were 35.23 (0.17), 35.68 (0.12), and 36.51 (0.17) °C with increasing  $T_w$ . Heat flux was measured at two sites on the skin adjacent to the temperature probe; mean values ( $\pm$  SE) were 87.7 (7.9), 71.8 (5.7), and 46.4 (2.8) W/m<sup>2</sup> with increasing  $T_w$ . Blood flows ( $\pm$  SE) entering the forearm and measured by plethysmography were 467 (82), 727 (86), and 779 (136) l/h/m<sup>3</sup> with increasing  $T_w$ .

**2. Analysis.** The approach taken herein to determine the net recovery of radial convective heat exchange was based on a simple heat balance. First, the convective heat exchange was obtained from the measured amount of heat lost through the skin after subtraction of the metabolic contribution assuming the Q<sub>10</sub> rule. Second, the potential convective heat loss (i.e., maximum possible) was determined by the difference between the average muscle temperature and the arterial blood temperature. One minus the ratio of these two quantities yields the fraction of heat recovery. All heat quantities are expressed as W/m<sup>3</sup> and apply to the muscle tissue only.

The convective heat exchange per unit volume ( $bc$ ) was determined by

$$bc = q_v - m$$

where  $q_v$  is the heat lost and  $m$  is the metabolic heat produced per unit volume. These quantities were determined by (3)

$$q_v = 2 \cdot hf / r$$

where  $hf$  and  $r$  are the heat flux and the radius at the muscle-fat boundary, and (4)

$$m = 684 \cdot 2^{(\bar{T} - 35)/10}$$

where  $\bar{T}$  is the average muscle temperature.

The potential convective heat loss (i.e. assuming no counter-current heat exchange) was determined by (4)

$$bc_{\max} = 1.13 \cdot bf \cdot (T_a - \bar{T})$$

where  $bf$  is the blood flow ( $l/h/m^3$ ) and  $T_a$  is the arterial blood temperature entering the forearm. Heat flux and blood flow at the muscle level were determined by assuming negligible metabolic heat production and blood flow at the skin+fat level.

## RESULTS

Values were calculated for each individual and means of these are reported below.

*Table of mean calculated values for forearm muscle tissue.*

$T_w$	$q_v$	$m$	$bc$	$bf$	$bc_{\max}$	$1 - bc/bc_{\max}$
15	5016	222	4794	467	9995	0.460
20	4076	306	3769	727	12032	0.678
30	2744	547	2197	779	5004	0.470

Convective heat exchange between blood and tissue accounts for most of the total heat produced (i.e.,  $bc + m$ ); the metabolic contribution is approximately 4, 7, and 20% of the total for  $T_w$  of 15, 20, and 30°C, respectively. Radial counter-current heat exchange returns about 53% of the potential convective heat available.

## CONCLUSION

Through the simple approach of comparing measured heat loss to the predicted potential heat loss using average values of tissue temperature and blood flow, the results suggest that radial counter-current heat exchange does not return all the heat brought in by the blood. This can be explained by noting that while heat recovery between counter-current vessels may be 100% efficient in the tissue space between the vessels, recovery is reduced in the much larger tissue space outside the conduction path between the vessels.

## REFERENCES

1. Weinbaum, S., Jiji, L.M. and Lemons, D.E., Theory and experiment for the vascular microstructure on surface tissue heat transfer - Part I: Anatomical foundation and model conceptualization. *J. Biomech. Eng.*, **107**: 131-139, 1984.
2. Ducharme, M.B., Vanhelder, W.P. and Radomski, M.W., Tissue temperature profile in the human forearm during thermal stress: I. Thermal stability. submitted
3. Sekins, K.M. and Emery, A.F., Thermal science for physical medicine. In *Therapeutic Heat and Cold*, edited by J.F. Lehmann, pp 97-99, Baltimore, MD, Williams and Wilkins, 1982.
4. Werner, J. and Buse, M., Temperature profiles with respect to inhomogeneity and geometry of the human body. *J. Appl. Physiol.* **65(3)**: 1110-1118, 1988.