

CORRELATION BETWEEN MEASURED SKIN WETTEDNESS AND SUBJECTIVE SENSATIONS OF SKIN WETNESS

G. Storaas¹ and M. K. Bakkevig²

Department of Zoology, Norwegian University of Science and Technology,
N-7055 Dragvoll

²SINTEF Unimed, N-7034 Trondheim, Norway

INTRODUCTION

The clothing insulation required to maintain thermal comfort during work in the cold will vary depending on the activity level. The insulation given by the clothing ensemble must be designed for the lowest work intensity and for rest periods when heat production is low. A thick clothing ensemble will reduce the dissipation of heat and water vapour by acting as a diffusion barrier and by absorbing sweat. During periods of high activity, the increased production of body heat and sweat may result in a sweat accumulation on the skin and in the underwear (Bakkevig and Nielsen, 1995). A correlation between skin wettedness and the feeling of discomfort has been claimed (Winslow et al. 1937). However, it is not clear whether we can sense increased levels of skin wettedness. This matter is addressed in this study which evaluates the correlation between measured skin wettedness and subjective sensations of skin wetness.

MATERIALS AND METHODS

Six male subjects, aged 24-29, participated in the experiments. The experiments were conducted in a climatic chamber ($T_a = +5^\circ\text{C}$) and consisted of two bouts of 40 minutes cycle exercise to ensure sweat production followed by 20 minutes rest after each period of exercise. Each subject conducted two tests where the work level in the exercise period was different, one approximated 40% of the subjects' $\dot{V}_{O_2, \text{max}}$ and the other 60% of $\dot{V}_{O_2, \text{max}}$. The experimental procedure with two work levels in the exercise period was chosen to ensure different levels of sweat production. A two layer clothing system was used in which the long-legged/long-sleeved underwear consisted of 80% wool and 20% polypropylene and the outerwear (trousers and jacket) consisted of 100% polypropylene. Relative humidity and temperature were measured at the skin at three locations (chest, back and thigh) during the experiments with the use of humidity sensors from VAISALA NMP 130 Y-series. They were placed between the skin and underwear with the aid of velcro. Local skin wettedness (w) on chest, back and thigh was calculated as




Figure 2 Correlation between measured skin wettedness on the chest and subjective evaluation of skin wettedness on the chest during work at $T_a=+5^{\circ}\text{C}$ $r=0.60$ ($n=24$).

DISCUSSION

The results, showing a correlation between measured skin wettedness and subjective ratings on general skin wetness, indicate that the test subjects were able to differentiate between different levels of skin wettedness. Whether this ability is a result of a perception of humidity per se or an indirect perception, is not clear. It has been shown (Withers P. C. 1990) that certain terrestrial insects have receptors in the skin that sense the water content of the surrounding air, so-called hygrometers. To our knowledge, this has never been found in mammals. The perception of skin wetness could be connected to the evaporative cooling of sweat on the skin. The efficiency of sweating as a cooling mechanism depends on allowing the moisture to evaporate from the skin and diffuse through the clothing to the environment in the form of water vapour. During work in a cold environment, a thick clothing ensemble will reduce the dissipation of heat and water vapour (Meinander, 1985). This will lead to a saturation of the water vapour concentration near the skin resulting in a low evaporation rate. After cessation of work, when heat and sweat production is reduced, the evaporation of accumulated sweat might increase. The rate of sweat evaporation during work might therefore differ, independently of the water vapour concentration and sweat accumulated, from the sweat evaporation during rest. As we found a