

# ANTHROPOMETRY AND PHYSICAL FITNESS IN MALE AND FEMALE SOLDIERS AND THEIR CAPABILITY TO CARRY LITTERS

W. von Restorff

Zentrales Institut des Sanitaetsdienstes der Bundeswehr  
Koblenz, Germany



## INTRODUCTION

For a variety of occupations, such as field medical specialists who have to transport stretcher patients without ergonomic aids in rough terrain, an evaluation of physical fitness prior to specialist training appears to be necessary. This may protect the employee from overburdening by and from excess demands by the employer.

Although there is some information available on the ability to carry loads (1,2) especially with respect to possible spine injuries (3), so far the data has not been correlated to more easily measurable predictors such as weight or stature (representing overall muscular strength) or isometric or isokinetic strength. Findings that hand grip endurance amounts, at best, to 15% of the maximum strength (1,4) have shed doubts on female capabilities of carrying litter patients (5). Comprehensive studies of female military fitness have been published (6,7). The data, however, cannot be applied to the Bundeswehr Medical Service, since selection and recruiting procedures might vary in addition to possible methodological or ethnic differences.

## METHODS

All 75 female and 60 self-selected male recruits of 4 different training cycles volunteered for the study. The measurements at the beginning and at the end of 10 weeks of basic military training (BMT) included weight, stature, sitting height, reaching height and body fat (skinfolds) (8,9). Isometric force was measured (baseline leg-chest-back dynamometer; Bückner, Sinzheim GE) in 4 positions: lifting from the squatting position; lifting from a height of 38 cm; lifting from the hip position; pressing upward from the shoulder position, together with right hand and left hand isometric grip force (Sadly Hand Dynamometer; Steeling Corp., Wood Dale, IL, USA). Carrying the patients was simulated either by sand bags weighing 60 to 90 kg on a stretcher (weighing 14 kg) or 2 water cans with equivalent weights for each hand. Successively increasing stretcher or water can loads had to be carried around an indoor volleyball court (55 m). Heart rate (Polar Elektro, Finland) was measured together with the time needed for the individual rounds. In some soldiers, O<sub>2</sub> consumption was monitored using a telemetry system (AERO-Sport, USA). All data was entered into data sheets and further processed using the MS Excel and SPSS PC+ statistics program. Statistically significant differences are seen at  $P < 0.05$ .

## RESULTS AND DISCUSSION

The study was designed as a "before and after test." Therefore, dropouts due to injury or other causes (e.g., other duty, retirement) could not be compensated. Thus, **only data** of recruits who participated in both measurements (55 females and 36 males) will be presented.

The anthropometric results are comparable to other published data (6,7). The changes of body weight, body fat amount and lean body mass of men and women during BMT were below 5% and statistically insignificant. This difference to previous studies (6) may be attributed to both the design of BMT and the short time period between the measurements.

The maximal isometric forces measured in the 4 different lift positions, both at the beginning and the end of BMT, are shown in Table 1, together with the

**Table 1.** Maximal isometric forces<sup>1</sup> at the start (S) and end (E) of BMT

<i>Men</i>	<u>Right hand</u>	<u>Left hand</u>	<u>Squat position</u>	<u>38-cm position</u>	<u>Standing position</u>	<u>Shoulder level</u>
Start of BMT <sup>2</sup>	549.9	513.4	1,447.0	1,608.3	944.5	1,547.8
	± 11.4	± 13.3	± 41.9	± 42.8	± 32.9	± 69.2
End of BMT <sup>2</sup>	581.0	534.4	1,544.5	1,587.0	1,046.1	1,612.9
	± 12.4	± 12.3	± 46.8	± 46.6	± 37.6	± 45.9
t-test, P <	0.066	0.245	0.120	0.735	0.044	0.429
S/E (%)	105.05	103.80	106.07	98.39	109.84	105.21
<i>Women</i>						
Start of BMT <sup>2</sup>	351.0	322.3	898.1	968.5	486.4	922.5
	± 6.5	± 6.7	± 20.7	± 25.3	± 10.3	± 33.7
End of BMT <sup>2</sup>	368.4	337.8	1,025.8	1,004.7	586.6	1,081.4
	± 6.6	± 7.1	± 22.6	± 25.7	± 16.3	± 37.0
t-test, P <	0.061	0.112	0.000	0.313	0.000	0.002
S/E (%)	105.0	104.8	114.2	103.7	120.6	117.2

<sup>1</sup>Force in Newtons. N = 36 male and 55 female recruits


<sup>2</sup>Values shown are means ± SEM.

maximal isometric handgrip strength values for men and women. The differences between men and women are statistically highly significant, which is not true, however, for **all** of the differences between beginning and end of BMT.

After BMT, women showed marked (> 10%) and statistically significant increases in strength during lift from the squatting, the standing positions and while pressing from the shoulder level position. These increases were less pronounced in men. **In** either group the slight differences of the handgrip strength readings did not reach the statistical significance level.

While all males were able to carry the simulated 90-kg patient at an average speed of 0.38 m/s, only 33 out of 75 female recruits were able to do so at the start of BMT (average speed 0.31 m/s). At the end of BMT, the number had improved to 29 out of 55 women. This improvement from 35% to 53% can be attributed to

the general conditioning during **BMT**. Heart rates rose during the carrying test to  $168 \pm 15$  or  $181 \pm 11$  bpm (males, females, respectively; means **SEM**). After BMT, the heart rates in both groups were lower; in males, however, the difference did not reach statistical significance. In females, the time needed to complete the carrying task was significantly reduced. During stretcher carrying, the  $O_2$  consumption approached  $> 25 \text{ ml} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$  equivalent to 83% of maximal  $VO_2$  in the females and documents the high workload posed on the young women.



**Figure 1.** Correlation analysis of speed while carrying a simulated 80-kg patient as determined by isometric grip strength of the stronger (max) or weaker (min) hands in 55 female recruits prior to and after BMT (start, end, respectively).

Out of a correlation matrix prepared to evaluate various possible predictors, the handgrip strength showed the strongest correlation with load carrying and transport speeds. The correlation turned out to be much stronger in females than in males, while there was no difference between the values of the respective stronger or weaker hand (max, min in Figure 1) or of the time of measurement.

## CONCLUSIONS

Measurements of various isometric force values revealed that the handgrip strength was the best predictor for the capacity to carry a simulated 90-kg patient. It is, therefore, suggested to administer a hand strength test prior to field medical specialist training in order to channel unsuited candidates to other career fields. Alternatively, special carrying training may be administered in order to prevent overburdening the young women.

## REFERENCES

1. Kilbom, A., Hägg, G.M. and Käll, C. 1992, One-handed load carrying—cardiovascular, muscular and subjective indices of endurance and fatigue, *European Journal of Applied Physiology*, 65, 52-58.
2. Laubach, L.L. 1979, Comparative muscular strength of men and women: a review of the literature, *Aviation, Space, and Environmental Medicine*, 47, 534-542.
3. Waters, T.R., Putz-Anderson, V., Garg, A. and Fine, L.J. 1993, Revised NIOSH equation for design and evaluation of manual lifting tasks, *Ergonomics*, 36,749-776
4. Schulze, H. 1993, ~~Zur~~ Leistungsfähigkeit von Finger- und Armmuskeln bei isometrischer Arbeit. Unpublished dissertation, Mainz.
5. von Restorff, W. and Land, K. 1994, Physical fitness in female recruits in the MC, in *Proceedings XXX International Congress of Military Medicine*, (Augsburg, Germany:ICMM).
6. Brock, J.R. and Legg, S.J. 1997, The effects of 6 weeks training on the physical fitness of female recruits to British army, *Ergonomics*, 40,400-411.
7. Teves, M.A., Wright, J.E. and Vogel, J.A. 1985, Performance on selected candidate screening test procedures before and after Army Basic and Advanced Individual Training, Technical Report No. T13/85, U.S. Army Research Institute for Environmental Medicine, Natick, MA.
8. Berres, F., Ulmer, H.V. and Lamberty, M. 1980, Calculation of total body fat from skinfold thickness by using an age corrected formula, *Pflugers Archiv—European Journal of Physiology*, 384, R 35.
9. Durnin, J.V.G.A. and Womersley, J. 1974, Body fat assessed from total body density and its estimation from skinfold thickness measurements on 481 men and women aged from 16 to 72 years, *British Journal of Nutrition*, 32, 77-97.