

WOMEN LITTER CARRIERS: OBSERVATION AND MODELING

W.R. Santee and W.T. Matthew

U.S. Army Research Institute of Environmental Medicine
Natick, MA USA



INTRODUCTION

Combat support roles for female soldiers expose them to extremes of both environment and physical exertion. We investigated tasks related to the evacuation of casualties by litter. Our goals were to quantify female performance under extreme conditions and to compare the efficacy of using male-based models to predict female performance.

MATERIALS AND METHODS

Data were collected on **4** female soldiers in a hot-dry environment (1). Investigators adhered to AR **70-25** and USAMRDC Regulation **70-25** on Use of Volunteers in Research. Subjects participated in these studies after giving their free and informed voluntary consent. Test activities consisted of walking (**1.34** m·s⁻¹) forward to the casualty and self-paced 2-person litter carriage (**68** kg load) to simulate stages in casualty evacuation. Metabolic costs for rest, walking and litter carriage were measured prior to testing. Litter carriage was intermittent work consisting of a series of short carries separated by rest periods. Clothing consisted of the Battledress Uniform (BDU) or the Battledress Overgarment (BDO). The BDU provided no chemical protection (MOPP-0 condition), whereas the BDO was worn with the M-40 mask (MOPP-4 condition). Data for rectal temperature (T_{re}) were compared to values calculated with the Heat **Strain** Decision Aid (HSDA) (2) and SCENARIO (3) models. The HSDA model is an executable version of the USAREM Heat Strain Model (4) developed to predict soldier performance. Model inputs are air temperature, relative humidity, wind speed, solar radiation, clothing, activity or metabolic rate and values for soldier height and weight and heat acclimation. The SCENARIO model was developed as a post hoc model to evaluate experimental results and requires additional inputs of maximum oxygen uptake and globe or mean radiant temperature. Although SCENARIO allows direct input of time-indexed metabolic rates, the HSDA does not. Consequently, metabolic inputs for the HSDA consisted of weighted average values that included a time adjustment to compensate for the metabolic cost of each lift.

RESULTS

Table 1 presents mean meteorological data used as modeling inputs, and Table 2 summarizes the test results. Mean values for subject height and weight were respectively 162 ± 1 cm and 63 ± 5 kg. There were statistically significant differences between MOPP-0 and MOPP-4 for ΔT_{re} ($P = 0.006$) while walking, but not for litter carriage. The primary limiting factor for litter carriage was muscular-skeletal stress.

Table 1. Environmental conditions during walking (W) and litter carriage (L) test activities in MOPP-0 (BDU) or MOPP-4 (BDO) clothing

Test	Meteorological values (SD)						
	T_a °C	T_g °C	wind $m \cdot s^{-1}$	RH %	WBGT °C	T_{rad} °C	T_{grnd} °C
W-BDU	39.41 (0.43)	5 1.11 (0.96)	2.49 (0.98)	12.57 (1.00)	28.9 (0.5)	82.20 (6.87)	52.95 (1.30)
W-BDO	37.96 (0.27)	47.05 (0.44)	4.82 (1.25)	17.85 (0.52)	28.3 (0.3)	84.05 (4.72)	50.52 (0.21)
LBDU	37.82 (0.34)	46.42 (0.65)	4.63 (1.13)	17.72 (1.57)	28.1 (0.4)	81.12 (4.86)	49.80 (1.17)
LBDO	35.46 (0.33)	42.21 (1.98)	4.82 (1.10)	26.79 (1.10)	27.4 (0.8)	71.53 (7.30)	46.38 (1.60)

Table 2. Summary of subject responses by activity¹

Test	T_{re} , °C	ΔT_{re} , °C	Endurance time in minutes range	<i>mean</i>
W-BDU	38.33 (0.42)	0.93 (0.27)	80-160 ²	140 (40)
W-BDO	38.55 (0.26)	1.28 (0.40)	20-37	29 (8)
LBDU	37.86 (0.50)	0.69 (0.51)	28-107	59 (35)
L-BDO	38.00 ³ (0.50)	0.76³ (0.41)	22-57³	433 (19)

¹ Values in parentheses are Standard Deviations

² Maximum time for any activity was 160 min

³ Three subjects, otherwise n=4

Figures 1 and 2 plot the **outputs** from both models with mean T_{re} values. The Root Mean Squared Deviations (RMSD) (**5**) from the observed T_{re} values were compared to the mean standard deviations (SD). For all days and models (4x2), with one exception, $RMSD \leq SD$. Overall, the HSDA model fit the **data** better. For litter carriage, the 2-day average RMSD for HSDA was 0.06 vs. 0.10 for **SCENARIO**.

DISCUSSION

With the exception of walking with chemical protection (**MOPP-4**), T_{re} values remained below 38.5°C. **This** indicates that factors other than thermal strain were limiting performance. Figures 1 and 2 show that both models project the general response patterns, although T_{re} is clearly overpredicted for walking in MOPP-4. When HSDA was used to model male performance during another **study** (**6**), the rate of increase in core temperature was overpredicted. It **may** be

desirable for a predictive model to provide a conservative estimate of core temperature, and thus provide some margin for error. The comparison of modeling results to this limited data set suggests there may be no such margin for error when the model is run with inputs for a female population.




Figure 1. Comparison of observed mean T_{re} to modeling results for walking in MOPP-0 (A) and MOPP-4 (B)




Figure 2. Comparison of observed mean T_{re} to modeling results for litter carriage in MOPP-0 (A) and MOPP-4 (B)

CONCLUSIONS

The results indicate that strength and equipment are limiting factors for litter carriage regardless of chemical protection (CP) clothing or weather condition. Without CP clothing, thermal strain was not the immediate limiting factor. For modeling, an accurate estimate of metabolic cost of each activity is critical. Both male-based models were applicable to our female subject population. The HSDA demonstrated better fit, but the format of SCENARIO is more suitable for direct input of metabolic rates for intermittent work.

REFERENCES

1. Santee, W.R., Kraning, **K.K.** and Matthew, W.T. **1997**, Modeling analysis of female litter bearers during heat stress, *Aviation, Space and Environmental Medicine*, **68**,647.
2. Joppa, **M.D.** **1993**, *P2NBC2 Heat Strain DecisionAid: User's Guide Version 2.1.*, (San Diego, CA:Scientific Applications International Corporation).
3. **Kraning, K.K.** and Gonzalez, R.R. **1997**, A mechanistic computer simulation of human work in heat that accounts for physical and physiological effects of clothing, aerobic fitness, and progressive dehydration, *Journal of Thermal Biology*, **22**, 331-342.
4. Pandolf, **K.B.**, Stroschein, L.A., Drolet, L.L., Gonzalez, R.R. and Sawka, M.N. **1986**, Prediction modeling of physiological responses and human performance in heat, *Computers in Biology and Medicine*, **16**, 310-329.
5. Haslam **R.A.** and Parsons **K.C.** **1989**, Models of human response to hot and cold environments, Final Report Vol. 1, Report No: APRE2170/106, Army Personnel Research Establishment, Farnborough, UK.
6. Santee, W.R., Matthew, **W.T.** and **Tharion**, W.J. **1992**, Simulated approach marches during thermal stress: A P2NBC2 study, Technical Report T12-92, U.S. Army Research Institute of Environmental Medicine, Natick, MA.

ACKNOWLEDGEMENT AND DISCLAIMER

The study was funded by the Defense Women's Health Research program. The views expressed in this report are those of the authors and should not be construed as official Department of the Army position, policy or decision, unless so designated by other official documentation.