

Differentiation

$y = f(x)$	$\frac{dy}{dx} = f'(x)$
k , constant	0
x	1
x^2	$2x$
x^3	$3x^2$
x^n , any constant n	nx^{n-1}
e^x	e^x
e^{kx}	ke^{kx}
$\ln kx = \log_e kx$	$\frac{1}{x}$

The sum-difference rule

$$\frac{d}{dx}(u(x) \pm v(x)) = \frac{du}{dx} \pm \frac{dv}{dx}$$

Constant multiples

$$\frac{d}{dx}(k \times f(x)) = k \times \frac{df}{dx}$$

for k constant

The product and quotient rules

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx} \quad \frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

The chain rule for differentiation

If $y = y(u)$ where $u = u(x)$ then $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$

Integration

$f(x)$	$\int f(x) dx$
k , constant	$kx + c$
x	$\frac{x^2}{2} + c$
x^2	$\frac{x^3}{3} + c$
x^n , ($n \neq -1$)	$\frac{x^{n+1}}{n+1} + c$
$x^{-1} = \frac{1}{x}$	$\ln x + c$
e^x	$e^x + c$
e^{kx}	$\frac{e^{kx}}{k} + c$


Written by Tony Croft, Geoff Simpson, and Mark Holmes
for the Mathematics Learning Support Centre
at Loughborough University

Typesetting and artwork by the authors

© 1997


Graphs of common functions

Linear: $y = mx + c$, m = gradient, c = vertical intercept




Exponential functions:

$e \approx 2.718$ is the exponential constant.



Quadratic functions: $y = ax^2 + bx + c$



(1) $b^2 - 4ac < 0$

(2) $b^2 - 4ac = 0$


(3) $b^2 - 4ac > 0$

(1) $b^2 - 4ac > 0$

(2) $b^2 - 4ac = 0$

(3) $b^2 - 4ac < 0$

Logarithmic functions:



Graphs of $y = \ln x$ and $y = \log_{10} x$

For the help you need to support your course

Economics Facts & Formulae

mathcentre is a project offering students and staff free resources to support the transition from school mathematics to university mathematics.

Arithmetic

When multiplying or dividing positive and negative numbers the sign of the result is given by:

positive \times positive = positive positive \times negative = negative
 negative \times positive = negative negative \times negative = positive

$$\frac{\text{positive}}{\text{positive}} = \text{positive} \quad \frac{\text{positive}}{\text{negative}} = \text{negative}$$

$$\frac{\text{negative}}{\text{positive}} = \text{negative} \quad \frac{\text{negative}}{\text{negative}} = \text{positive}$$

The BODMAS rule reminds us of the order in which operations are carried out. BODMAS stands for:

Brackets () First priority
Of \times Second priority
Division \div Second priority
Multiplication \times Second priority

Addition + Third priority
Subtraction - Third priority

Fractions.

$$\text{fraction} = \frac{\text{numerator}}{\text{denominator}}$$

Adding and subtracting fractions. To add or subtract two fractions first rewrite each fraction so that they have the same denominator. Then, the numerators are added or subtracted as appropriate and the result is divided by the common denominator: e.g.

$$\frac{4}{5} + \frac{3}{4} = \frac{16}{20} + \frac{15}{20} = \frac{31}{20}$$

Multiplying fractions. To multiply two fractions, multiply their numerators and then multiply their denominators:

e.g.

$$\frac{3}{7} \times \frac{5}{11} = \frac{15}{77}$$

Dividing fractions. To divide two fractions, invert the second and then multiply: e.g.

$$\frac{3}{5} \div \frac{2}{3} = \frac{3}{5} \times \frac{3}{2} = \frac{9}{10}$$

Algebra

Removing brackets:

$$a(b + c) = ab + ac, \quad a(b - c) = ab - ac$$

$$(a + b)(c + d) = ac + ad + bc + bd$$

Formula for solving a quadratic equation:

$$\text{if } ax^2 + bx + c = 0 \text{ then } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Laws of Indices:

$$a^m a^n = a^{m+n} \quad \frac{a^m}{a^n} = a^{m-n} \quad (a^m)^n = a^{mn}$$

$$a^0 = 1 \quad a^{-m} = \frac{1}{a^m} \quad a^{1/n} = \sqrt[n]{a} \quad a^{\frac{m}{n}} = (\sqrt[n]{a})^m$$

Laws of Logarithms:

$y = \log_b x$ means $b^y = x$ and b is called the **base**.
 e.g. $\log_{10} 2 = 0.3010$ means $10^{0.3010} = 2.000$, to 4 s.f.

$$\log_b AB = \log_b A + \log_b B, \quad \log_b \frac{A}{B} = \log_b A - \log_b B,$$

$$\log_b A^n = n \log_b A, \quad \log_b 1 = 0, \quad \log_b b = 1$$

Logarithms to base e , denoted \log_e or alternatively \ln are called *natural logarithms*. The letter e stands for the exponential constant which is approximately 2.718.

Proportion and Percentage

To convert a fraction to a percentage multiply by 100 and label the result as a percentage.

Examples

$$\frac{5}{8} \text{ as a percentage is } \frac{5}{8} \times 100\% = 62.5\%$$

$$\frac{1}{3} \text{ as a percentage is } \frac{1}{3} \times 100\% = 33\frac{1}{3}\%$$

Some common conversions are

$$\frac{1}{10} = 10\%, \quad \frac{1}{4} = 25\%, \quad \frac{1}{2} = 50\%, \quad \frac{3}{4} = 75\%$$

Ratios are simply an alternative way of expressing fractions. Consider dividing £200 between two people in the ratio of 3:2. This means that for every £3 the first person gets, the second person gets £2. So the first gets $\frac{3}{5}$ of the total, and the second gets $\frac{2}{5}$ of the total; that is £120 and £80.

Generally, to split a quantity in the ratio $m : n$, the quantity is divided into $\frac{m}{m+n}$ of the total and $\frac{n}{m+n}$ of the total.

Sigma notation

The Greek capital letter sigma, Σ , is used as an abbreviation for an addition sum. Suppose we have n values x_1, x_2, \dots, x_n and we wish to add them together. The sum

$$x_1 + x_2 + \dots + x_n \text{ is written } \sum_{i=1}^n x_i$$

Note that i runs through all whole number values from 1 to n . So, for instance

$$\sum_{i=1}^3 x_i \text{ means } x_1 + x_2 + x_3$$

Example

$$\sum_{i=1}^5 i^2 \text{ means } 1^2 + 2^2 + 3^2 + 4^2 + 5^2$$

Statistics

Population values, or **parameters**, are denoted by Greek letters. Population mean = μ . Population variance = σ^2 . Population standard deviation = σ . Sample values, or **estimates**, are denoted by roman letters.

The **mean** of a sample of n observations x_1, x_2, \dots, x_n is

$$\bar{x} = \frac{\sum_{i=1}^n x_i}{n} = \frac{x_1 + x_2 + \dots + x_n}{n}$$

The unbiased estimate of the **variance** of these n sample observations is

$$s^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n-1} \text{ which can be written as}$$

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n x_i^2 - \frac{n\bar{x}^2}{n-1}$$

The sample unbiased estimate of **standard deviation**, s , is the square root of the variance:

$$s = \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n-1}}$$

The Greek alphabet

A	α	alpha	I	ι	iota	P	ρ	rho
B	β	beta	K	κ	kappa	Σ	σ	sigma
Γ	γ	gamma	Λ	λ	lambda	T	τ	tau
Δ	δ	delta	M	μ	mu	Υ	υ	upsilon
E	ϵ	epsilon	N	ν	nu	Φ	ϕ	phi
Z	ζ	zeta	Ξ	ξ	xi	X	χ	chi
H	η	eta	O	o	omicron	Ψ	ψ	psi
Θ	θ	theta	Π	π	pi	Ω	ω	omega