

RESPONSE TO EXERCISE AMONG MALE AND FEMALE ARMY OFFICERS

Ulf Bergh, Ulf Danielsson and Christer Weinö

National Defence Research Establishment, 17290 Stockholm

War Academy, 10425 Stockholm

INTRODUCTION

Women are entering areas such as military and police forces where most people are males. To enter such jobs, tests of physical work capacity have to be passed (e. g. running or cycle ergometry). The question is often: should tests and requirements be identical for males and females. One possible reason for having different requirements would be if the tests did not take into account gender-specific characteristics of importance for the job. For example, tests are often of short duration while jobs might require considerable endurance and therefore such tests will probably not account for all factors of importance for job performance. This study was undertaken to explore if a given task, performed in a given environment, using standard equipment caused different response in male compared to female officers.

METHODS

The subjects were nine male (age 26 ± 1 years, height 184 ± 6 cm and weight 82 ± 9 kg) and eight female (age 27 ± 2 years, height 170 ± 4 cm and weight 63 ± 6 kg) army officers. There were no significant difference between males and females either in age or running time for 2 km and 10 km (males 7.8 ± 0.2 min and 47.8 ± 2.9 min; females 8.2 ± 0.3 min and 49.2 ± 2.5 min). The task consisted of marching on an indoor course for 2 • 50 min with a 10 min rest in between. The course contained sharp turns and obstacles that elevated the centre of gravity by 1.45 m per lap. Each lap was 92 m long and was covered in 1 min which gave a walking speed of 1.53 m/s (5.5 km/h). The air temperature and relative humidity were 20 °C and 40%, respectively. The subjects were dressed in combat uniform and 22 kg of equipment including an armoured vest. The day before, as well as 30 min after the march, cycle ergometry was performed consisting of 6 min of submaximal exercise at a work rate proportional to body mass raised to 2/3 power, followed by 6 min of maximal exercise, where the subjects were instructed to perform as many pedal revolutions as possible within these 6 min. Average power output was used as a performance criterion and was denoted P_{\max} . Body mass (m_b), mass of body + uniform + equipment (m_{bdr}), rectal temperature (T_{re}), haemoglobin concentration (Hb), and haematocrit (Hct) were measured before and after the march. Heart rate (HR) was monitored once every minute. Metabolic rate (MR) was measured, with a portable device, during the first 10 min of either the first or the second half of the march. Every 10 min the subjects stopped for 15-20 s while the mass of body plus equipment was monitored. At the same time the subjects rated their perceived exertion (RPE), temperature (RPT) and comfort (RPC).

RESULTS AND DISCUSSION

Both males and females had higher physical work capacity than the average person of corresponding sex and age, the females being considerably more extreme in this respect. Hence, present data does not represent a valid comparison between average males and females. One should also keep in mind that all of the subjects had passed selection processes before and during their military training.

The males performed significantly better than the females at the maximal cycle test (307 W vs 251 W). During the march, males displayed significantly greater body mass loss (Δm_b), evaporation, MR, and significantly lower HR compared to females (see table 1). P_{\max} was lower after the march than before. There was no gender

difference in $\Delta m_b/A_d$, $\Delta m_{bdr}/A_d$ and $\Delta m_b/m_b$, meaning that compensating for differences in body surface area (A_d) eliminated all differences observed between genders. This is different from the findings of Sawka et al. (1983) who found that women had significantly lower sweat rates per unit of surface area. There were, however, difference in procedures between these studies, e.g. in the study of Sawka et al the subjects were very lightly dressed and was allowed to drink ad libitum. The magnitude of changes induced by the march in HR, RPE, RPT, RPC, Hb, Hct and P_{max} were not significantly different between genders. Thus, the physical strain of the march did not seem to be different for males as compared to females. The differences observed between genders most likely originated from different body *size*, mainly body mass. Therefore, the advantage of a lower fluid loss found and lower metabolic rate observed among the female officers should not be interpreted as a gender-specific characteristic.

Table 1. Mean values (M) and standard deviations (s) for some of the variables measured in connection with the march.

VARIABLE	MALES		FEMALES	
	M	s	M	s
Heart rate (beats \cdot min ⁻¹)	138	16	155	7
Metabolic rate (W)	657	38	536	64
Rectal temperature (°C)	38,3	0,4	38,4	0,3
Δ body mass (kg)	1,720	0,227	1,237	0,283
Amass of body + uniform + equip. (kg)	0,939	0,082	0,795	0,159
$(\Delta m_b/m_b) \cdot 100$ (%)	2,11	0,33	1,95	0,34
Δ HR march (beats \cdot min ⁻¹)	14	7	17	6
Δ Tre march (°C)	0,8	0,5	0,8	0,2
Δ Haematokrit (%)	-0,3	34	-1	3,3

CONCLUSIONS

The differences found between genders were explained primarily by differences in body mass and therefore this study does not support the use of different requirements for male and females in tests of physical endurance capacity.

REFERENCE

Sawka M., M. Toner, R. Francesconi, and K. Pandolf. 1983. Hypohydration and exercise: effects of heat acclimation, gender, and environment. *J. Appl. Physiol.: Respirat. Environ. Exercise Physiol.* 55(4): 1147-1153

ACKNOWLEDGEMENTS

The authors are indebted to J. Keinanen, B. Ståhl, G. Larsson, A. Mikiver, M. Mikiver, E. Malm, and S. Grambo for skilful technical assistance.