

THE EFFECTS OF EXERCISE, HEAT AND MICROCLIMATE COOLING ON TERMINAL STROKE VOLUME IN MEN AND WOMEN

J.H. Heaney¹, M.J. Buono² & J.A. Hodgdon¹

¹Naval Health Research Center, San Diego, CA, USA

²San Diego State University, San Diego, CA, USA



INTRODUCTION

We have previously reported that in lieu of the more common end-of-test criteria such as core temperature $> 39^{\circ}\text{C}$, and heart rate (HR) > 180 bpm (2,3), volitional fatigue for men exercising in the heat appears to occur with a reduction in stroke volume (SV) to a value of approximately 75 ml/beat/min (1). Since it has been established that females have a lower absolute SV (4) it is doubtful that a terminal stroke volume (TSV) of 75 ml/beat/min would be observed in females. It was hypothesized that using either body surface area (BSA), which would account for an increased peripheral blood circulation and the exchange between the skin and the environment, or fat free mass, which represents the body mass to be perfused would provide a physiological metric to normalize SV results between males and females. Additionally, we observed that while the use of a microclimate cooling garment (ice vest) attenuated the rate of cardiovascular drift, the fall in terminal SV (TSV) was independent of temperature and ice vest usage for males exercising in the heat. This paper provides results from a comparison evaluation of male and female SV responses to exercising in the heat both with and without the use of microclimate cooling,

MATERIAL AND METHODS

This research was conducted in compliance with all Federal regulations concerning the protection of human subjects. Fourteen males (24 ± 5 yrs; 1.94 ± 0.1 m²; 17.3 ± 4.5 % fat; 49.2 ± 4.5 ml·kg⁻¹·min⁻¹ VO_{2max}) and seventeen females (24 ± 4 yrs; 1.71 ± 0.1 m²; 30.6 ± 4.8 % fat; 43.9 ± 6.5 ml·kg⁻¹·min⁻¹ VO_{2max}) provided written consent. Upon completing an eight day acclimation protocol each subject underwent duplicate heat exposures trials once wearing an ice vest (V) and once without (NV) in three different thermal environments (A = 43°C and 48% RH; B = 51°C and 33% RH; and C = 57°C and 25% RH) for a total of six heat exposure trials. During each heat trial the subjects alternated approximately 20 min of treadmill walking (3mph, 3% grade) with 40 min of seated rest for six hours or until volitional fatigue. The microclimate cooling garment for this study was the Steele ice vest which contains six gel packs placed in three horizontal pouches across the chest and back region. The ice packs were frozen at -29°C and replaced every 120min in environment A and every 90 min in environments B and C. The ice vest has a total weight of 5.1kg and a total cooling capacity of 270 watt·hr⁻¹. Heart rate (HR), and rectal temperature (T_{re}), were

measured every min throughout the heat trial. Cardiac output (CO) values, determined using the indirect Fick CO₂ rebreathing technique, were assessed approximately 7-10 min into each exercise session and provided a SV value. TSV results were evaluated as absolute values, adjusted for BSA (TSV/BSA), and adjusted for lean body mass (TSV/FFM). Menstrual phase was not controlled, however post-hoc blood analysis revealed that female subject participation during the follicular and luteal phase was evenly distributed.

RESULTS

Since there were several subjects who were able to complete the full 6-hours in the A-environment ice vest condition (mean = 5.5 hours), the physiological results from this trial did not represent a volitional fatigue effort and therefore, were not included in the analysis. End-of-test values for T_{re} (°C), HR (bpm), CO (L·min⁻¹), TSV (ml·beat⁻¹), TSV/BSA (ml·beat⁻¹·m⁻²), TSV/FFM (ml·beat⁻¹·kg⁻¹), and stay time (min) were analyzed for effects of gender using a repeated measures ANOVA. Physiological results for the 5 trials are displayed in Table 1 (males) and Table 2 (females).

Table 1. Male Physiological Results.

n = 14	<u>A No-Vest</u>	<u>B No-Vest</u>	<u>B IceVest</u>	<u>C No-Vest</u>	<u>C IceVest</u>
* T_{re}	38.7f0.5	38.8f0.5	37.8f0.4	38.9f0.4	38.4f0.5
*HR	149 ± 12	147 ± 13	134 ± 13	153 ± 15	147f 15
*CO	11.5f1.6	10.3f1.8	10.6f1.5	11.2f1.7	11.0f1.9
*TSV	78f14	71 ± 16	80 f 14	74f 13	75f 16
*TSV/BSA	40f6	37 f 7	41 f 7	33 ± 7	39 ± 8
TSV/FFM	1.22f0.2	1.10 ± 0.2	1.24f0.2	1.16 ± 0.2	1.17f0.2
*Stay time	162f 32	139 ± 24	340 ± 35	38 ± 22	242f 62

* = significant gender effect, $p < 0.05$.

Table 2. Female physiological Results.

n = 17	<u>A No-Vest</u>	<u>B No-Vest</u>	<u>B IceVest</u>	<u>C No-Vest</u>	<u>C IceVest</u>
* T_{re}	38.9k0.4	38.9f0.4	38.5f0.5	38.9 ± 0.4	38.9f0.5
*HR	156 ± 17	162 ± 15	159 ± 15	161 ± 20	162 ± 14
*CO	8.9f2.2	8.5 ± 1.4	9.3f1.0	9.0f2.0	9.0f1.7
*TSV	55 ± 13	53 f 10	59 f 7	57 ± 15	56f 12
*TSV/BSA	32 ± 7	31 I 5	34 ± 4	33 ± 8	33 ± 6
TSV/FFM	1.19f0.3	1.13f0.2	1.27f0.2	1.21f0.3	1.21 ± 0.2
*Stay time	134 ± 37	93 f 26	240 I 92	71 ± 6	168f 68

* = significant gender effect $p < 0.05$.

Use of the ice vest resulted in stay times over twice as long as stay times with no-vest. In general, end-of-test T_{re} values were similar for males and females, however, end-of-test HR values for females were 13 bpm greater. As expected, higher CO and SV values were observed for males and are most likely associated with their greater body size compared to females. The absolute TSV values were significantly different ($p < 0.05$) between males and females. In fact male TSV values ($76 \text{ ml}\cdot\text{beat}^{-1}$) were similar to the first SV (SV1) female values (73 debeat^{-1}). While the absolute SV1 and end-of-test or TSV values were significantly different ($p < 0.05$), the qualitative response in SV was almost identical. The average SV1 value for females was 32% lower (108 vs. 73, male and female, respectively) while the average TSV value for females was 26% lower ($76 \text{ ml}\cdot\text{beat}^{-1}$ vs. $56 \text{ ml}\cdot\text{beat}^{-1}$, male and female, respectively). Since BSA was thought to significantly effect the circulatory response (via increased peripheral blood pooling) and therefore impact the drop in SV, absolute TSV values were adjusted for BSA. Although this adjustment attenuated the TSV difference ($39 \text{ ml}\cdot\text{beat}^{-1}\cdot\text{m}^{-2}$ vs. $33 \text{ ml}\cdot\text{beat}^{-1}\cdot\text{m}^{-2}$, for males and females, respectively), female TSV values were still significantly lower ($P < 0.05$). In an attempt to further explain the TSV gender differences, TSV was next adjusted for FFM. The males in this study had a mean FFM of 66 kg vs. 46 kg for the female participants. Figure 1 shows that adjusting TSV for FFM eliminated all TSV gender differences ($P > 0.05$).




Figure 1. Absolute Terminal Stroke Volume and FFM Adjusted Terminal Stroke Volume Results

With respect to environmental, cooling, and gender conditions, Figure 1 supports the hypothesis that TSV is a valid indicator of volitional fatigue. Despite end-of-test T_{re} values $< 39^{\circ}\text{C}$, and end-of-test HR values < 162 bpm, and stay times that ranged from 71 min to 340 min, the participants in this study terminated their heat exposure trial when TSV/FFM fell to approximately $1.2 \text{ ml}\cdot\text{beat}^{-1}\cdot\text{kg}^{-1}$.

CONCLUSIONS

Absolute terminal stroke volume was significantly higher in males. Adjusting terminal stroke volume for BSA reduced the magnitude of the gender differences in stroke volume but did not eliminate it. However, terminal stroke volume was identical for males and females when adjusted for fat free mass. Wearing an ice vest significantly increased stay times by more than twice the stay times without the vest and significantly delayed the fall in stroke volume. These results suggest that the limiting factor in maintaining an adequate stroke volume may be more closely associated with perfusion of the body mass as a whole rather than peripheral blood circulation at the skin surface.

REFERENCES

1. Heaney, J.H., Buono, M.J., Wilmore, K.M., and Banta, G.R. 1992, Heart rate, core temperature, and stroke volume as indicators of volitional fatigue during exercise in the heat. *Proceedings of the Fifth International Conference on Environmental Ergonomics*, Maastricht, The Netherlands, 14-15.
2. Armstrong, L.E., Hubbard, R.W., Deluca, J.P., and Christensen, E.L. 1987, Signs and symptoms of heat exhaustion during strenuous exercise. *Annals of Sports Medicine*, Vol 3, pp 182-189.
3. Dassler, A.R. 1977, Heat stress, work function and physiological heat exposure limits in man. In: *Thermal Analysis-Human Comfort-Indoor Environments*, National Bureau of Standards, NBS Special Publication #491, pp 65-92.
4. Jones, N.L. and Campbell, E.J. *Clinical Exercise Testing*, 1982, W.B. Sanders Company, Philadelphia PA, pp. 250-251.